

Niatab™ 500

Pure Sustained-Release, High Potency Niacin

DESCRIPTION

Niatab™ 500 tablets, provided by Douglas Laboratories®, slowly release 500 mg of pure niacin. Sustained release niacin is better tolerated than regular crystalline high-potency niacin supplements.

FUNCTIONS

Niacin (vitamin B3) occurs in the body as two metabolically active coenzymes, NAD (nicotinamide adenine dinucleotide) and NADP (NAD phosphate).

The niacin coenzymes NAD and NADP have pervasive roles in energy-related and biosynthetic metabolic processes. At least 200 enzymes depend on these niacin cofactors. The NAD-dependent enzymes are involved in mostly catabolic, oxidative reactions that release energy from carbohydrate, fat, and protein, whereas the NADP-dependent enzymes more commonly function in biosynthetic pathways of such compounds as fatty acids and steroid hormones.

Independent of its functions as NAD or NADP, niacin is also involved in the regulation of normal blood lipoprotein and cholesterol levels. † Dietary niacin is generally well absorbed and taken up by the liver which converts it to NAD and NADP. Any excess niacin is metabolized by the liver and excreted by the kidneys. Because the liver has a limited capacity to metabolize niacin, high doses of regular, crystalline niacin are often not well tolerated, causing vasodilatory side effects known as skin-flushing and itching. For maximum benefit and fewer side effects, Niatab™ 500 releases its niacin over an extended period of time. This way, the niacin is more efficiently utilized by the liver and more evenly distributed to peripheral tissues compared to crystalline niacin. For the same reasons, Niatab™ 500 can generally be used at somewhat lower doses than crystalline niacin supplements.

INDICATIONS

Niatab™ 500 tablets may be a useful nutritional adjunct for individuals who wish to increase their intake of niacin without the uncomfortable side effects of skin flushing.

FORMULA (#NTB)

Tablet:

Niacin..... 500 mg

SUGGESTED USE

One-half to one tablet, one to two times daily during or after a meal (i.e., breakfast and dinner), or as directed by physician. As a general guideline when substituting Niatab™ 500 for crystalline niacin, use Niatab™ 500 at 1/2 to 2/3 of the level used for crystalline niacin.

SIDE EFFECTS

No adverse side effects have been reported.

STORAGE

Store in a cool, dry place, away from direct light. Keep out of reach of children.

REFERENCES

Alderman JD et al. Am J Cardiol 1989;64:725-729.

Niatab™ 500
Pure Sustained-Release, High Potency Niacin

Canner PL et al. J Am Coll Cardiol 1986;8:1245-1255.

Colletti RB et al. Pediatrics 1993;92:78-82.

Keenan JM et al. Arch Intern Med 1991;151:1424-1432.

Keenan J, et al. Archives Of Internal Medicine [serial online]. July 1991;151(7):1424-1432.

Lavie CJ et al. Am J Cardiol 1992;69:1083-1085.

Probstfield JL. Arch Int Med 1994;154:1557-1559.

For more information visit douglaslabs.com

† These statements have not been evaluated by the Food and Drug Administration.
This product is not intended to diagnose, treat, cure, or prevent any disease.

Manufactured by
Douglas Laboratories
600 Boyce Road
Pittsburgh, PA 15205
800-245-4440
douglaslabs.com



© 2015 Douglas Laboratories. All Rights Reserved